



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

93. Proposed by JOHN R. JEFFREY, Student in Ohio State University, Columbus, Ohio.

Solve the following differential equation :

$$(1-x^2) \frac{d^2 y}{dx^2} - x \frac{dy}{dx} + y = 2x, \text{ when } x < 1.$$

94. Proposed by ALOIS F. KOVARIK, Instructor in Mathematics and Physics, Decorah Institute, Decorah, Ia.

Find the minimum isosceles triangle that can be described about a given ellipse, having its base parallel to the major axis. [Ex. 16, page 166, *Rice and Johnson's Differential Calculus*.

*** Solutions of these problems should be sent to J. M. Colaw not later than August 10.

MECHANICS.

91. Proposed by CHARLES C. CROSS, Whaleyville, Va.

The bow of a boat which is a inches wide is inclined at an angle α . When in motion in perfectly calm water the water was found to rise b inches on the bow. Required the velocity of the boat.

92. Proposed by WALTER H. DRANE, Graduate Student, Harvard University, Cambridge, Mass.

A particle, starting at the vertex, slides down a smooth parabolic curve. Prove that in order to leave the curve at the extremity of the latus rectum, the initial velocity of the particle must be $pg[1/\sqrt{2}-1]$ where p is semi-latus rectum.

*** Solutions of these problems should be sent to B. F. Finkel not later than August 10.

AVERAGE AND PROBABILITY.

75. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy, Irving College, Mechanicsburg, Pa.

Find the mean area of all plane rectilinear right triangles having a constant perimeter p .

76. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy, Irving College, Mechanicsburg, Pa.

In a given ellipse, the extremities of a focal chord are joined with the center. Find the average area of the triangle thus formed.

*** Solutions of these problems should be sent to B. F. Finkel not later than August 10.

MISCELLANEOUS.

79. Proposed by S. HART WRIGHT, A. M., Ph. D., Penn Yan, N. Y.

In latitude $42^\circ 30' N. = \lambda$, a tree 100 feet long $= \alpha$, leans in the direction $S. 60^\circ W. = \beta$, with an angle of elevation with the level ground, of $30^\circ = \gamma$. The sun's declination being $1^\circ 36' 24'' N. = \delta$, in what direction will the shadow of the tree point, when the sun is on the meridian?